APPROVED USE

LOKELMA is a prescription medication for the treatment of high levels of potassium in the blood (hyperkalemia) in adults. LOKELMA should not be used as an emergency treatment for life-threatening hyperkalemia because of its delayed onset of action

It's going to be

LOKELMA can help

Worried about high potassium?

Hyperkalemia — high potassium — can lead to serious health issues if left untreated.

Read on to **find out how LOKELMA can help lower potassium levels and keep them low with continued use.** Plus, learn how to take LOKELMA, and what to expect during treatment.

IMPORTANT SAFETY INFORMATION FOR LOKELMA® (sodium zirconium cyclosilicate)

What is the most important thing I should know about LOKELMA?

Don't take LOKELMA if you have:

- severe constipation, a blockage (obstruction) in your bowel, or hard stool that will not pass (impaction)
- problems with your bowel movements after surgery

LOKELMA may not work and may worsen gastrointestinal conditions.

LOKELMA®
(sodium zirconium cyclosilicate)
5 g | 10 g for oral suspension

Please see Important Safety Information on page 6, along with the full Prescribing Information for LOKELMA.

What is high potassium, and what does it mean for you?

A higher than normal level of potassium (K^+) in the blood, also known as hyperkalemia, can be an ongoing and serious condition if left untreated. High potassium may cause complications, such as heart issues, that can lead to hospitalization.



mmol/L=millimole per liter.

*In some cases, up to 5.2 mmol/L is considered normal.

†Like high potassium, low potassium can also potentially cause serious health issues.

Who is at risk?

Some people may have a higher chance of developing hyperkalemia, such as those with:

- Chronic Kidney Disease (CKD)
- Heart failure
- High blood pressure
- Diabetes
- Certain medications to treat these conditions

IMPORTANT SAFETY INFORMATION FOR LOKELMA (Cont'd)

LOKELMA®
(sodium zirconium cyclosilicate)
5 g | 10 g for oral suspension

Please see Important Safety Information on page 6, along with the full Prescribing Information for LOKELMA.

What is the most important thing I should know about LOKELMA? (continued)

LOKELMA can cause fluid retention (edema) in your feet, ankles, hands or other parts of the body.

- LOKELMA contains 400 mg of sodium in each 5 g dose. Watch for signs of fluid retention (edema) in your body especially if you limit sodium in your diet or are likely to have fluid retention in your body due to heart or kidney problems. Your doctor may tell you

Why LOKELMA?

LOKELMA lowers your high potassium level and helps keep it in a normal range with continued use.

- It's designed to **target and remove** potassium
- It has **no taste or smell**
- **It had few side effects**

In fact, LOKELMA was generally well tolerated — with few side effects — in studies involving more than 1700 people who had hyperkalemia and were not on dialysis.

In clinical studies with people with high potassium who are not on dialysis, **LOKELMA started to bring potassium levels down in as early as 1 hour after the first dose. Plus, it helped keep potassium levels normal with continued use.** In these studies, patients were not required to follow a low potassium diet.

LOKELMA should not be used as an emergency treatment for life-threatening hyperkalemia because of its delayed onset of action.

How does LOKELMA work?

LOKELMA is a potassium binder that works in the digestive tract, where it binds potassium and removes it from the body so it doesn't enter the bloodstream.

LOKELMA is designed to target and remove potassium and not calcium and magnesium.

LOKELMA contains approximately 400 mg of sodium in each 5-g dose, but the extent of absorption by the patient is unknown.

High potassium can be an ongoing but treatable condition. It is important to keep taking LOKELMA as prescribed.

IMPORTANT SAFETY INFORMATION FOR LOKELMA (Cont'd)

What is the most important thing I should know about LOKELMA? (continued)

to reduce the sodium in your diet or to increase the dose of your diuretic as needed

If you are a hemodialysis patient:

- Do not take LOKELMA on dialysis days
- Tell your doctor if you suddenly feel sick such as decreased oral intake of food or fluids or have diarrhea, as you may be at risk of developing low levels of potassium in your blood while on LOKELMA; therefore, your dose of LOKELMA may need to be adjusted

Tell your doctor(s) you are taking LOKELMA before having an abdominal X-ray as it may appear on the X-ray.

Please see Important Safety Information on page 6, along with the full Prescribing Information for LOKELMA.

Taking LOKELMA

LOKELMA is a white to grey powder that you mix with water and drink. Don't worry — it has no taste or smell. When you fill your prescription, you will receive LOKELMA in individual, foil-lined packets. Continue to take LOKELMA exactly as prescribed for as long as your doctor prescribes it.

How to take LOKELMA

1. Fill your cup with at least 3 tablespoons of water

2. Empty the entire contents of the LOKELMA packet(s) into the cup

3. Stir well and drink immediately. If there is powder left, add more water, stir, and drink again. Repeat until all the powder is gone

LOKELMA can be taken with or without food and does not need to be refrigerated

2 Hrs.

If you have been prescribed other oral medications, **you may need to take LOKELMA 2 hours before or 2 hours after your medication**. Talk to your doctor about all other medications you are currently taking

Your doctor is here to help. Ask if you need to make any changes to your medication schedule.

LOKELMA®
(sodium zirconium cyclosilicate)
5 g | 10 g for oral suspension

Please see Important Safety Information on page 6, along with the full Prescribing Information for LOKELMA.

IMPORTANT SAFETY INFORMATION FOR LOKELMA (Cont'd)

What are the most common side effects with LOKELMA?

The most common side effect of LOKELMA is swelling in your body caused by fluid retention (edema). Tell your doctor if you have any side effects that bother you or do not go away.

For patients who are not on dialysis, LOKELMA can help maintain normal potassium levels with continued use

Initial treatment for people not on dialysis

TAKE ONE
10-g PACKET **3X**/DAY
FOR UP TO 48 HOURS

Maintenance treatment for people not on dialysis

TAKE ONE
10-g PACKET **1X**/DAY
Your doctor may monitor your potassium level to help find the dose that is right for you.

For patients who are receiving dialysis, LOKELMA can help keep potassium levels within a normal range between dialysis sessions

Recommended dose for people on dialysis

TAKE ONE
5-g PACKET **1X**/DAY
ON NON-DIALYSIS DAYS ONLY
Your doctor may monitor your potassium level to help find the dose that is right for you.

IMPORTANT SAFETY INFORMATION FOR LOKELMA (Cont'd)

What are the most common side effects with LOKELMA? (continued)

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements. LOKELMA may affect how some oral medicines work. Ask your doctor or pharmacist if you need to take LOKELMA 2 hours before or after other oral medicines.

Please see Important Safety Information on page 6, along with the full Prescribing Information for LOKELMA.

Important Safety Information

What is the most important thing I should know about LOKELMA?
Don't take LOKELMA if you have:

- severe constipation, a blockage (obstruction) in your bowel, or hard stool that will not pass (impaction)
- problems with your bowel movements after surgery

LOKELMA may not work and may worsen gastrointestinal conditions.

LOKELMA can cause fluid retention (edema) in your feet, ankles, hands or other parts of the body.

- LOKELMA contains 400 mg of sodium in each 5 g dose. Watch for signs of fluid retention (edema) in your body especially if you limit sodium in your diet or are likely to have fluid retention in your body due to heart or kidney problems. Your doctor may tell you to reduce the sodium in your diet or to increase the dose of your diuretic as needed

If you are a hemodialysis patient:

- Do not take LOKELMA on dialysis days
- Tell your doctor if you suddenly feel sick such as decreased oral intake of food or fluids or have diarrhea, as you may be at risk of developing low levels of potassium in your blood while on LOKELMA; therefore, your dose of LOKELMA may need to be adjusted

Tell your doctor(s) you are taking LOKELMA before having an abdominal X-ray as it may appear on the X-ray.

What are the most common side effects with LOKELMA?
The most common side effect of LOKELMA is swelling in your body caused by fluid retention (edema). Tell your doctor if you have any side effects that bother you or do not go away.

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements. LOKELMA may affect how some oral medicines work. Ask your doctor or pharmacist if you need to take LOKELMA 2 hours before or after other oral medicines.

APPROVED USE
LOKELMA is a prescription medication for the treatment of high levels of potassium in the blood (hyperkalemia) in adults.

LOKELMA should not be used as an emergency treatment for life-threatening hyperkalemia because of its delayed onset of action.

Please see US Full Prescribing Information for LOKELMA.

LOKELMA®
(sodium zirconium cyclosilicate)
5 g | 10 g for oral suspension

You may report side effects related to AstraZeneca products.

Finding support

It's going to be OK — let our support team do some of the lifting

My LOKELMA Support is dedicated to helping you better understand your high potassium and overall wellness. **You'll receive financial and insurance-related guidance, as well as a direct-mail welcome kit** to ease you into your treatment with LOKELMA.

You'll have access to:

- Personal insurance coverage counselors
- Educational resources
- Information about savings

My LOKELMA Support Program

Ways to sign up

Call 1-844-LOKELMA (1-844-565-3562)
Monday - Friday, 8 AM to 8 PM EST

Visit join-lokelma.com

Pay as little as $0 a month* with the LOKELMA savings card!

Download and bring the $0 Co-pay Savings Card and a valid prescription to your pharmacy (no activation required).

You can download your LOKELMA Savings Card at lokelma-savings.com

*Subject to eligibility rules; restrictions apply.

Visit lokelma.com to learn more

LOKELMA®
(sodium zirconium cyclosilicate)
5 g | 10 g for oral suspension

Please see Important Safety Information on page 6, along with the full Prescribing Information for LOKELMA.

LOKELMA is a registered trademark of the AstraZeneca group of companies.
©2026 AstraZeneca. All rights reserved. US-113221 Last Updated 6/26